

**EFICO Alpha 717**


**TROUT**

**Declaration**

**3 - 4,5 - 6 mm**

Crude protein	42,0 %
Crude lipid	22,0 %
Carbohydrates (NFE)	17,5 %
Fibre	3,5 %
Ash	6,0 %
Total phosphorus (P)	1,1 %
Gross energy (MJ/kcal)	21,6 / 5174
Digestible energy (MJ/kcal)	18,4 / 4408


**Energy distribution**



**Composition**

- Fishmeal
- Soya meal
- Blood meal
- Fish oil
- Wheat
- Rapeseed cake
- Sunflower meal
- Field beans
- Rape seed oil
- Vitamins - Minerals

**Ecological value**



Due to natural variations in the raw materials, the information about carbohydrates, fibre, and ash, and the composition may vary compared to the stated values. See label for further information.

**Feeding guides (kg feed per 100 kg fish per day)**

Lowest possible feed conversion rate – to be used when optimal feed utilisation is required

Fish size g	Fish size cm	Pellet size mm	2°C	4°C	6°C	8°C	10°C	12°C	14°C	16°C	18°C	20°C
50 - 100	16 - 20	3,0	0,88	1,03	1,18	1,35	1,51	1,67	1,80	1,87	1,86	1,70
100 - 200	20 - 26	4,5	0,78	0,91	1,04	1,19	1,33	1,47	1,59	1,66	1,64	1,50
200 - 300	26 - 29	4,5	0,69	0,80	0,92	1,05	1,18	1,30	1,40	1,46	1,45	1,32
300 - 450	29 - 33	4,5	0,62	0,71	0,82	0,94	1,05	1,16	1,26	1,31	1,30	1,18
450 - 600	33 - 36	6,0	0,57	0,66	0,77	0,87	0,98	1,08	1,17	1,22	1,21	1,10
600 - 800	36 - 40	6,0	0,51	0,59	0,68	0,78	0,87	0,96	1,04	1,08	1,08	0,98
800 - 1000	40 - 43	6,0	0,50	0,58	0,67	0,77	0,86	0,95	1,02	1,07	1,06	0,97

Optimal feeding – to be used when an optimal relation between large production and good feed utilisation is required

Fish size g	Fish size cm	Pellet size mm	2°C	4°C	6°C	8°C	10°C	12°C	14°C	16°C	18°C	20°C
50 - 100	16 - 20	3,0	0,88	1,03	1,23	1,64	2,15	2,73	3,31	3,70	3,50	1,86
100 - 200	20 - 26	4,5	0,78	0,91	1,07	1,43	1,88	2,39	2,89	3,23	3,06	1,62
200 - 300	26 - 29	4,5	0,69	0,80	0,93	1,25	1,64	2,09	2,53	2,83	2,68	1,42
300 - 450	29 - 33	4,5	0,62	0,71	0,83	1,12	1,47	1,87	2,27	2,54	2,40	1,27
450 - 600	33 - 36	6,0	0,57	0,66	0,77	1,04	1,36	1,74	2,11	2,36	2,23	1,18
600 - 800	36 - 40	6,0	0,51	0,59	0,69	0,93	1,22	1,55	1,89	2,11	1,99	1,05
800 - 1000	40 - 43	6,0	0,50	0,58	0,69	0,92	1,21	1,54	1,87	2,09	1,98	1,05

Feeding should be adapted to the chosen production strategy and current farming conditions.